

| Salt Point State Park & Kruse Rhododendron State Natural Reserve- Vascular Plants - (version 2019-08) | | | | |
|---|---|-----------------------|--|-------------------|
| α | Botanical Name | Common Name | Habitat | Life History/Form |
| FERNS | | | | |
| Athyriaceae - Lady Fern | | | | |
| | <i>Athyrium filix-femina</i> var. <i>cyclosorum</i> | lady fern | moist woods, forests | perennial |
| Blechnaceae (Deer Fern Family) - 2 taxa | | | | |
| | <i>Struthiopteris spicant</i> | deer fern | moist woods, canyons | perennial |
| | <i>Woodwardia fimbriata</i> | giant chain fern | along creeks, in springs, seeps | perennial |
| Cystopteridaceae - Fragile Fern | | | | |
| | <i>Cystopteris fragilis</i> | fragile fern | moist, shaded streambanks | perennial |
| Dennstaedtiaceae (Bracken Family) - 1 taxon | | | | |
| | <i>Pteridium aquilinum</i> var. <i>pubescens</i> | bracken | widespread in grassland, scrub | perennial |
| Dryopteridaceae (Wood Fern Family) - 2 taxa (see also Woodsiaeeae) | | | | |
| | <i>Dryopteris arguta</i> | wood fern | woods, forests | perennial |
| | <i>Polystichum munitum</i> | western sword fern | damp forests, scrub, along streams | perennial |
| Equisetaceae (Horsetail Family) - 3 taxa | | | | |
| | <i>Equisetum arvense</i> | common horsetail | wet soils near streams, seeps | perennial |
| | <i>Equisetum hyemale</i> ssp. <i>affine</i> | common scouring rush | moist scrub near streams | perennial |
| | <i>Equisetum telmateia</i> ssp. <i>braunii</i> | giant horsetail | streambanks, wet soils | perennial |
| Ophioglossaceae (Adder's-tongue Family) - 1 taxon | | | | |
| | <i>Sceptridium multifidum</i> | grape-fern; moonwort | wet grassland, scrub | perennial |
| Polypodiaceae (Polypody Family) - 3 taxa | | | | |
| | <i>Polypodium californicum</i> | California polypody | damp woods, stream banks | perennial |
| | <i>Polypodium glycyrrhiza</i> | licorice fern | damp canyons, on rocks, logs | perennial |
| | <i>Polypodium scouleri</i> | leather-leaf fern | on trees, rocks, soil in salt spray zone | perennial |
| Pteridaceae (Brake Family) - 3 taxa | | | | |
| | <i>Adiantum aleuticum</i> | five-finger fern | wooded riparian | perennial |
| | <i>Adiantum jordanii</i> | California maidenhair | damp woods | perennial |
| | <i>Pentagramma triangularis</i> | goldback fern | banks, cliffs, logs | perennial |
| GYMNOSPERMS | | | | |
| Cupressaceae (Cypress Family) - 3 taxa | | | | |
| 1B.2 | <i>Hesperocyparis pygmaea</i> | pygmy cypress | pygmy forest | small tree |
| 1B.2 | <i>Hesperocyparis macrocarpa</i> * | Monterey cypress | coastal grassland | tree |

| | | | |
|---|-------------------------|-------------------------------|-----------------------------------|
| <i>Sequoia sempervirens</i> | coast redwood | redwood forest | tree |
| Pinaceae (Pine Family) - 6 taxa | | | |
| <i>Abies grandis</i> | grand fir | forest | tree |
| <i>Pinus muricata</i> | Bishop pine | forest | tree |
| <i>Pinus radiata</i> * | Monterey pine | disturbed sites; forest | tree |
| <i>Pseudotsuga menziesii</i> | Douglas-fir | forest | tree |
| <i>Tsuga heterophylla</i> | western hemlock | forest | tree |
| Taxaceae (Yew Family) - 1 taxon | | | |
| <i>Torreya californica</i> | California nutmeg | moist, shady woodland, forest | shrub; tree |
| NYMPHAEALES | | | |
| Nymphaeaceae (Water-lily Family) - 1 taxon | | | |
| <i>Nuphar polysepala</i> | yellow pond-lily | open, still water | perennial |
| MAGNOLIIDS | | | |
| Aristolochiaceae (Pipevine Family) - 1 taxon | | | |
| <i>Asarum caudatum</i> | wild-ginger | streamsides; moist thickets | perennial |
| Lauraceae (Laurel Family) - 1 taxon | | | |
| <i>Umbellularia californica</i> | California bay | woodland; forest | shrub; tree |
| EUDICOTS | | | |
| Adoxaceae - 1 taxon | | | |
| <i>Sambucus racemosa</i> var. <i>racemosa</i> | red elderberry | wet places; scrub | shrub |
| Aizoaceae (Carpetweed or Fig-marigold Family) - 3 taxa | | | |
| <i>Carpobrotus chilensis</i> * | sea fig | | perennial (trailing, mat-forming) |
| <i>Carpobrotus edulis</i> * | Hottentot-fig; iceplant | | perennial (trailing, mat-forming) |
| <i>Tetragonia tetragonoides</i> * | New Zealand spinach | disturbed sites | annual |
| Amaranthaceae (Amaranth Family) - 2 taxa | | | |
| <i>Chenopodium californicum</i> | California goosefoot | sandy places (Stump Beach) | perennial |
| <i>Extriplex californica</i> | California orach | sandy places; marshes | perennial |
| Anacardiaceae (Cashew or Sumac Family) - 1 taxon | | | |
| <i>Toxicodendron diversilobum</i> | poison-oak | scrub; woodland; forest | shrub; vine |
| Apiaceae (Carrot, Parsley, Dill Family) aka Umbelliferae - 20 taxa | | | |
| <i>Angelica hendersonii</i> | coast angelica | scrub; grassland | perennial |
| <i>Apium graveolens</i> * | celery | damp areas | annual; perennial |
| <i>Conioselinum pacificum</i> | conioselinum | scrub | perennial |

| | | | | |
|---|---|-----------------------|--|-------------------------|
| | <i>Conium maculatum</i> * | poison hemlock | moist, disturbed sites; scrub, streamsides | perennial |
| | <i>Daucus carota</i> * | Queen Anne's lace | grassland; disturbed sites | annual |
| | <i>Daucus pusillus</i> | rattlesnake weed | grassland | annual |
| | <i>Eryngium armatum</i> | coyote thistle | moist grassland | perennial |
| | <i>Foeniculum vulgare</i> * | fennel | disturbed sites; roadsides | perennial |
| 4.2 | <i>Glehnia littoralis</i> ssp. <i>leiocarpa</i> | beach silvertop | coastal strand | perennial |
| | <i>Heracleum maximum</i> | cow parsnip | moist coastal scrub, grassland | perennial |
| | <i>Ligusticum apiifolium</i> | celery-leaved lovage | shady banks, grassland | perennial |
| | <i>Oenanthe sarmentosa</i> | ditch-carrot | marshes; streamsides | perennial |
| | <i>Osmorhiza berteroii</i> | sweet cicely | moist, shady woodland, forest | perennial |
| 4.2 | <i>Perideridia gairdnerii</i> ssp. <i>gairdnereii</i> | Gairdner yampah | grassland; woodland | perennial |
| | <i>Sanicula arctopoides</i> | footsteps-of-spring | coastal bluffs; grassland | perennial |
| | <i>Sanicula bipinnatifida</i> | purple sanicle | grassland | perennial |
| | <i>Sanicula crassicaulis</i> | Pacific sanicle | grassland; woodland | perennial |
| | <i>Sanicula laciniata</i> | cutleaf sanicle | grassland; scrub | perennial |
| | <i>Scandix pecten-veneris</i> * | Venus's needle | disturbed sites; grassland | annual |
| | <i>Torilis arvensis</i> * | tall sock-destroyer | woodland; scrub; grassland | annual |
| Apocynaceae (Dogbane Family) - 1 taxon | | | | |
| | <i>Vinca major</i> * | periwinkle | disturbed sites; homesteads | perennial (groundcover) |
| Araliaceae (Ginseng Family) 2 taxa | | | | |
| | <i>Aralia californica</i> | elk-clover; spikenard | moist sites; streamsides | perennial |
| | <i>Hedera helix</i> * | English ivy | forest; disturbed sites | perennial (vine) |
| Asteraceae (Sunflower Family) aka Compositae - 71 taxa | | | | |
| | <i>Achillea millefolium</i> | yarrow | grassland | perennial |
| | <i>Adenocaulon bicolor</i> | trailfinder | shaded woodland, forest | perennial |
| | <i>Agoseris apargioides</i> | seaside agoseris | sandy soils; grassland; dunes | perennial |
| | <i>Agoseris heterophylla</i> | annual agoseris | grassland | annual |
| | <i>Ambrosia chamissonis</i> | beach-bur | beach; dunes | perennial |
| | <i>Anaphalis margaritacea</i> | pearly everlasting | grassland; scrub | perennial |
| | <i>Anisocarpus madioides</i> | woodland madia | woodland; forest | perennial |
| | <i>Arnica discoidea</i> | rayless arnica | woodland | perennial |
| | <i>Artemisia californica</i> | California sagebrush | scrub; grassland | shrub |
| | <i>Artemisia douglasiana</i> | Douglas mugwort | wetlands; grassland; scrub | perennial |
| | <i>Artemisia suksdorfii</i> | Suksdorf mugwort | wetlands; scrub; grassland | perennial |

| | | | |
|--|--------------------------------------|--|----------------------|
| <i>Baccharis pilularis</i> var. <i>consanguinea</i> | coyote brush | scrub; grassland | shrub |
| <i>Bellis perennis</i> * | English daisy | moist grassland; disturbed sites | perennial |
| <i>Calycadenia multiglandulosa</i> | sticky rosin weed | dry grassland, slopes | annual |
| <i>Carduus pycnocephalus</i> ssp. <i>p.</i> * | Italian thistle | grassland; disturbed sites | annual |
| <i>Cichorium intybus</i> * | chicory | disturbed sites; grassland | perennial |
| <i>Cirsium brevistylum</i> | short-styled thistle | stream corridors; disturbed sites | biennial |
| <i>Cirsium douglasii</i> var. <i>breweri</i> | swamp thistle | wet grassland; scrub | biennial; perennial |
| <i>Cirsium occidentale</i> var. <i>occidentale</i> | cobweb thistle | grassland; coastal slopes; dunes | biennial |
| <i>Cirsium quercetorum</i> | brownie thistle | grassland | perennial |
| <i>Cirsium vulgare</i> * | bull thistle | grassland; scrub; disturbed, moist sites | biennial |
| <i>Corethrodyne filaginifolia</i> | sand-aster | grassland; slopes | perennial |
| <i>Cotula coronopifolia</i> * | brass-buttons | coastal wetlands | perennial |
| <i>Deinandra corymbosa</i> ssp. <i>corymbosa</i> | tarplant | grassland | annual |
| <i>Delairea odorata</i> * | Cape-ivy | scrub; woodland; disturbed sites | perennial (vinelike) |
| <i>Erigeron canadensis</i> | Canadian horseweed | disturbed sites | annual |
| <i>Erigeron glaucus</i> | seaside daisy | coastal bluffs, grassland | perennial |
| <i>Eriophyllum lanatum</i> var. <i>arachnoideum</i> | woolly sunflower | banks, slopes; scrub | perennial |
| <i>Eriophyllum staechadifolium</i> | lizardtail; seaside woolly sunflower | coastal bluffs; scrub | shrub |
| <i>Euchiton gymnocephalus</i> * | creeping cudweed | forest flats, along trails | |
| <i>Eurybia radulina</i> | rough-leaved aster | woodland; scrub | perennial |
| <i>Gamochaeta ustulata</i> | purple cudweed | grassland; disturbed sites | annual; perennial |
| <i>Grindelia stricta</i> var. <i>platyphylla</i> | gumplant | grassland; coastal bluffs, slopes | perennial |
| <i>Helenium puberulum</i> | sneezeweed | wetlands | annual; perennial |
| <i>Hemizonella minima</i> | little madia | woodland; forest; scrub | annual |
| <i>Hemizonia congesta</i> ssp. <i>clevelandii</i> | hayfield tarweed | grassland (upper prairie) | annual |
| <i>Hemizonia congesta</i> ssp. <i>lutescens</i> | pale yellow hayfield tarplant | coastal prairie | annual |
| 1B.2 <i>Hesperevax sparsiflora</i> var. <i>brevifolia</i> | short-leaved revax | coastal bluffs; rocky sites | annual |
| <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i> | few-flowered evax | grassland; scrub | annual |
| <i>Heterotheca sessiliflora</i> ssp. <i>bolanderi</i> | Bolander golden-aster | grassland; coastal bluffs | perennial |
| <i>Hieracium albiflorum</i> | hawkweed | dry woodland; forest | perennial |
| <i>Hypochaeris glabra</i> * | smooth cat's-ear | grassland; dunes; disturbed sites | annual |
| <i>Hypochaeris radicata</i> * | rough cat's-ear | grassland; disturbed sites | perennial |
| 1B.2 <i>Lasthenia californica</i> ssp. <i>macrantha</i> | perennial goldfields | grassland | perennial |

| | | | |
|---|-----------------------------|-----------------------------------|-------------------|
| <i>Lasthenia glaberrima</i> | marsh lasthenia | salt marsh; wetlands | annual |
| <i>Lasthenia gracilis</i> | common goldfields | grassland | annual |
| <i>Layia gaillardioides</i> | layia | scrub; grassland | annual |
| <i>Layia platyglossa</i> ssp. <i>platyglossa</i> | tidy-tips | grassland; coastal bluffs | annual |
| <i>Leontodon saxatilis</i> * | hawkbit | grassland | annual; perennial |
| <i>Leucanthemum vulgare</i> * | ox-eye daisy | coastal strand | perennial |
| <i>Logfia filaginoides</i> | daggerleaf cottonrose | grassland; scrub | annual |
| <i>Madia exigua</i> | threadstem madia | grassland; woodland | annual |
| <i>Madia sativa</i> | coast madia | grassland; scrub | annual |
| <i>Matricaria discoidea</i> * | pineapple weed | disturbed sites | annual |
| <i>Microseris laciniata</i> | cut-leaf microseris | grassland | perennial |
| <i>Petasites frigidus</i> var. <i>palmatus</i> | coltsfoot | streamsides; shaded wetlands | perennial |
| <i>Pseudognaphalium californicum</i> | California cudweed | scrub; grassland | annual; biennial |
| <i>Pseudognaphalium luteoalbum</i> * | everlasting cudweed | grassland; disturbed sites | annual |
| <i>Pseudognaphalium ramosissimum</i> | pink cudweed | grassland; scrub | biennial |
| <i>Pseudognaphalium stramineum</i> | Chilean cudweed | scrub; slopes | annual; biennial |
| <i>Psilocarphus tenellus</i> var. <i>tenellus</i> | woolly-heads | disturbed, compacted soil | annual |
| <i>Senecio glomerata</i> * | Australian fireweed | scrub, grassland, woodland | perennial |
| <i>Senecio minima</i> * | Australian fireweed | scrub, grassland, woodland | annual; perennial |
| <i>Senecio vulgaris</i> * | groundsel | disturbed sites | annual |
| <i>Silybum marianum</i> * | milk thistle | disturbed sites | annual; biennial |
| <i>Solidago elongata</i> | West Coast Canada goldenrod | scrub | perennial |
| <i>Solidago spathulata</i> | coast goldenrod | sandy soils; dunes | perennial |
| <i>Soliva sessilis</i> * | field burweed | disturbed, compacted soil | annual |
| <i>Sonchus asper</i> ssp. <i>asper</i> * | prickly sow-thistle | disturbed sites; grassland | annual |
| <i>Sonchus oleraceus</i> * | common sow-thistle | disturbed sites; grassland; scrub | annual |
| <i>Symphyotrichum chilense</i> | California coast aster | scrub; grassland | perennial |
| <i>Taraxacum officinale</i> * | dandelion | disturbed, moist sites; grassland | perennial |
| <i>Wyethia angustifolia</i> | narrow-leaved mule's-ears | grassland | perennial |

Berberidaceae (Barberry Family) - 2 taxa

| | | | |
|--------------------------------|--------------------------------|-------------------------|------------------------|
| <i>Berberis nervosa</i> | longleaf mahonia | woodland | shrub |
| <i>Vancouveria planipetala</i> | redwood-ivy; inside-out flower | moist coniferous forest | subshrub (groundcover) |

Betulaceae (Birch Family) - 2 taxa

| | | | |
|---------------------------------|-----------|-------------|------|
| <i>Alnus rubra</i> ¹ | red alder | streamsides | tree |
|---------------------------------|-----------|-------------|------|

| | | | |
|--|-----------------------------|-----------------------------------|-----------------------|
| <i>Corylus cornuta</i> var. <i>californica</i> | California hazel | damp woodland; forest | shrub |
| Boraginaceae (Borage Family) - 8 taxa (includes Hydrophyllaceae) | | | |
| <i>Amsinckia spectabilis</i> | seaside fiddleneck | coastal grassland; dunes | annual |
| <i>Myosotis discolor</i> * | blue scorpion-grass | disturbed sites | annual; biennial |
| <i>Nemophila menziesii</i> var. <i>atomaria</i> | baby white-eyes | grassland | annual |
| <i>Nemophila parviflora</i> var. <i>parviflora</i> | small-flowerednemophila | shaded places | annual |
| <i>Phacelia bolanderi</i> | Bolander phacelia | moist coastal bluffs | perennial |
| <i>Phacelia californica</i> | California phacelia | rocky places | perennial |
| <i>Phacelia malvifolia</i> | stinging phacelia | gravelly, sandy slopes in scrub | annual |
| <i>Plagiobothrys reticulatus</i> var. <i>reticulatus</i> | reticulate popcorn flower | wet grassland | annual |
| <i>Plagiobothrys undulatus</i> | wavy-stemmed popcorn flower | grassland | |
| Brassicaceae (Mustard Family) aka Cruciferae - 13 taxa | | | |
| <i>Barbarea vulgaris</i> * | common yellow cress | wet sites | perennial |
| <i>Brassica nigra</i> * | black mustard | disturbed sites | annual |
| <i>Brassica rapa</i> * | field mustard | disturbed sites, grassland | annual |
| <i>Cakile maritima</i> * | sea rocket | dunes, beaches | annual |
| <i>Capsella bursa-pastoris</i> * | shepherd's purse | disturbed sites | annual |
| <i>Cardamine californica</i> var. <i>californica</i> | milk maids | grassland, woodland, forest | perennial |
| <i>Cardamine oligosperma</i> | few-seeded bitter cress | disturbed sites, grassland, scrub | annual |
| 1B.2 <i>Erysimum concinnum</i> | bluff wallflower | grassland | bien/perennial |
| <i>Hirschfeldia incana</i> * | summer mustard | disturbed sites | biennial to perennial |
| <i>Lepidium didymum</i> * | swinecress | disturbed sites | annual |
| <i>Lepidium nitidum</i> var. <i>nitidum</i> | peppergrass | disturbed sites | annual |
| <i>Nasturtium officinale</i> | water cress | wet places, streams, seeps | perennial |
| <i>Raphanus sativus</i> * | wild radish | disturbed sites | annual |
| Campanulaceae (Bellflower Family) - 2 taxa | | | |
| <i>Asyneuma prenanthoides</i> | nodding harebell | forest slopes | perennial |
| 1B.2 <i>Campanula californica</i> | swamp harebell | freshwater swamp | perennial |
| Caprifoliaceae (Honeysuckle Family) - 3 taxa (see also Adoxaceae) | | | |
| <i>Lonicera hispidula</i> | honeysuckle | woodland; scrub | perennial vine |
| <i>Lonicera involucrata</i> var. <i>ledebourii</i> | twinberry | wet scrub; streamsides | shrub |
| <i>Symporicarpos albus</i> var. <i>laevigatus</i> | snowberry | wet places; streamsides | shrub |
| Caryophyllaceae (Pink Family) - 14 taxa | | | |
| <i>Cardionema ramosissimum</i> | sand mat | coastal grassland; dunes | perennial |

| | | | |
|--|----------------------|--|-------------------------|
| <i>Cerastium arvense</i> | field chickweed | grassland; rocky sites | perennial |
| <i>Cerastium glomeratum</i> * | mouse-ear chickweed | grassland; disturbed sites | annual |
| <i>Polycarpon tetraphyllum</i> * | four-leaved allseed | grassland; dunes; disturbed sites | annual |
| <i>Sagina apetala</i> | dwarf pearlwort | disturbed sites; grassland | annual |
| <i>Sagina maxima</i> ssp. <i>crassicaulis</i> | big pearlwort | coastal bluffs | perennial |
| <i>Sagina procumbens</i> | Arctic pearlwort | moist, sandy or gravelly places | perennial |
| <i>Silene gallica</i> * | windmill pink | grassland; disturbed sites | annual |
| <i>Spergula arvensis</i> * | stickwort | grassland; disturbed sites | annual |
| <i>Spergularia macrotheca</i> var. <i>macrotheca</i> | sticky sand-spurrey | rocky, sandy coastal grassland, bluffs | perennial |
| <i>Spergularia marina</i> | coastal sand-spurrey | muddy, alkaline marshy places | annual |
| <i>Spergularia rubra</i> * | sand-spurrey | disturbed, compacted soil | annual; perennial |
| <i>Stellaria media</i> * | common chickweed | moist, disturbed sites | annual |
| <i>Stellaria pallida</i> * | chickweed | coastal grassland | annual |
| Celastraceae (Staff-tree Family) - 2 taxa | | | |
| <i>Euonymus occidentalis</i> | western burning bush | moist, shady canyons | shrub |
| <i>Paxistima myrsinoides</i> | Oregon boxwood | shaded slopes in woodland | subshrub |
| Convolvulaceae (Morning-glory Family) - 6 taxa | | | |
| <i>Calystegia occidentalis</i> | morning-glory | scrub | perennial |
| <i>Calystegia purpurata</i> ssp. <i>purpurata</i> | coast morning-glory | grassland, scrub | perennial |
| 1B.2 <i>Calystegia purpurata</i> ssp. <i>saxicola</i> | morning-glory | scrub | perennial |
| <i>Calystegia soldanella</i> | beach morning-glory | coastal strand; dunes | perennial |
| <i>Convolvulus arvensis</i> * | field bindweed | upper prairie | perennial |
| <i>Cuscuta</i> sp. | dodder | on <i>Lupinus arboreus</i> | |
| <i>Dichondra donelliana</i> | dichondra | grassland | perennial (mat-forming) |
| Crassulaceae (Stonecrop Family) - 2 taxa | | | |
| <i>Crassula connata</i> | sand pygmy-weed | disturbed sites; grassland | annual |
| <i>Dudleya farinosa</i> | coast live-forever | coastal bluffs; rocky sites | perennial |
| Cucurbitaceae (Gourd Family) - 2 taxa | | | |
| <i>Marah fabacea</i> | coast man-root | scrub; grassland; woodland | perennial (trailing) |
| <i>Marah oregana</i> | Oregon man-root | scrub; canyons | perennial (trailing) |
| Dipsacaceae (Teasel Family) - 1 taxon | | | |
| <i>Dipsacus sativus</i> * | Fuller's teasel | disturbed sites; roadside banks | biennial |
| Ericaceae (Heather Family) - 16 taxa | | | |
| <i>Allotropa virgata</i> | sugar stick | forest, in humus | perennial (non-green) |

| | | | |
|--|----------------------|--|-----------------------|
| <i>Arbutus menziesii</i> | Pacific madrone | woodland; forest | tree |
| <i>Arctostaphylos columbiana</i> | hairy manzanita | woodland openings | shrub |
| <i>Arctostaphylos glandulosa</i> | Eastwood manzanita | pygmy forest/scrub | shrub |
| <i>Arctostaphylos nummularia</i> | shatterberry | woodland openings; pygmy forest | shrub |
| <i>Arctostaphylos uva-ursi</i> forma <i>coactilis</i> | bearberry | marine terrace | shrub (low) |
| <i>Chimaphila menziesii</i> | prince's pine | forest | perennial |
| <i>Chimaphila umbellata</i> | prince's pine | redwood-tanoak forest | perennial |
| <i>Gaultheria shallon</i> | salal | woodland; forest; scrub | shrub (low) |
| <i>Hemitomes congestum</i> | gnome plant | forest, in humus | perennial (non-green) |
| <i>Pleurocospora fimbriolata</i> | fringed pinesap | forest, in humus | perennial (non-green) |
| <i>Rhododendron columbianum</i> | western Labrador-tea | coastal wetlands | shrub |
| <i>Rhododendron macrophyllum</i> | California rose-bay | woodland; forest; pygmy forest | shrub |
| <i>Rhododendron occidentale</i> | western azalea | moist sites; streambanks | shrub |
| <i>Vaccinium ovatum</i> | blue huckleberry | woodland; forest; scrub | shrub |
| <i>Vaccinium parvifolium</i> | red huckleberry | woodland; forest | shrub |
| Euphorbiaceae (Spurge Family) - 2 taxa | | | |
| <i>Croton setiger</i> | turkey-mullein | disturbed sites, grassland | annual |
| <i>Euphorbia peplus</i> * | petty spurge | disturbed moist sites | annual |
| Fabaceae (Legume Family) aka Leguminosae - 41 taxa | | | |
| <i>Acacia pycnantha</i> * | golden wattle | disturbed sites (old homestead) | tree |
| <i>Acmispon americanus</i> var. <i>americanus</i> | Spanish clover | disturbed sites; grassland | annual |
| <i>Acmispon brachycarpus</i> | colchita | grassland; open, gravelly sites, slopes | annual |
| <i>Acmispon grandiflorus</i> var. <i>grandiflorus</i> | large-flowered lotus | pine, Douglas-fir forest openings | perennial |
| <i>Acmispon heermannii</i> var. <i>orbicularis</i> | Heermann lotus | scrub | perennial |
| <i>Acmispon junceus</i> var. <i>biolettii</i> | lotus | coastal sand; scrub | perennial |
| <i>Acmispon parviflorus</i> | miniature lotus | grassland; open sites, slopes | annual |
| <i>Acmispon wrangelianus</i> | Wrangel lotus | grassland | annual |
| <i>Astragalus nuttallii</i> var. <i>virgatus</i> | Nuttall milkvetch | sandy soils; coastal strand; scrub; bluffs | perennial |
| <i>Cytisus scoparius</i> * | Scotch broom | disturbed sites; roadsides; grassland | shrub |
| <i>Genista monspessulana</i> * | French broom | disturbed sites; roadsides; grassland | shrub |
| 4.2 <i>Hosackia gracilis</i> | harlequin lotus | moist grassland | perennial |
| | rose-flowered lotus | disturbed sites in forest | perennial |
| | hillside pea | scrub; woodland | perennial |
| | hairy annual lotus | disturbed sites; grassland | annual |

| | | | |
|--|----------------------|---------------------------------------|------------------|
| <i>Lotus corniculatus</i> * | bird's-foot trefoil | disturbed sites | perennial |
| <i>Lotus tenuis</i> * | bird's-foot trefoil | disturbed sites; grassland | perennial |
| <i>Lupinus arboreus</i> var. <i>arboreus</i> * | yellow bush lupine | scrub; dunes | shrub |
| <i>Lupinus variicolor</i> | vari-colored lupine | grassland; coastal bluffs | perennial |
| <i>Medicago polymorpha</i> * | California burclover | disturbed sites | annual |
| <i>Melilotus albus</i> * | white sweetclover | disturbed sites; roadsides | annual; biennial |
| <i>Melilotus indicus</i> * | sourclover | disturbed sites; roadsides | annual |
| <i>Trifolium angustifolium</i> * | narrow-leaved clover | roadsides | annual |
| <i>Trifolium barbigerum</i> | bearded clover | grassland | annual |
| <i>Trifolium dubium</i> * | little hop clover | disturbed sites; grassland | annual |
| <i>Trifolium fragiferum</i> * | strawberry clover | trails | perennial |
| <i>Trifolium hirtum</i> * | rose clover | roadsides | annual |
| <i>Trifolium incarnatum</i> * | crimson clover | roadsides | annual |
| <i>Trifolium macraei</i> | Chile clover | grassland; disturbed sites | annual |
| <i>Trifolium microcephalum</i> | smallhead clover | grassland; disturbed sites | annual |
| <i>Trifolium microdon</i> | squarehead clover | grassland; disturbed sites | annual |
| <i>Trifolium oliganthum</i> | few-flowered | moist grassland, disturbed sites | annual |
| <i>Trifolium repens</i> * | white clover | moist, disturbed sites | perennial |
| <i>Trifolium subterraneum</i> * | subterranean clover | grassland; disturbed sites | perennial |
| <i>Trifolium variegatum</i> | whitetip clover | grassland; moist sites | annual |
| <i>Trifolium willdenovii</i> | tomcat clover | grassland; scrub; open sites | annual |
| <i>Trifolium wormskiioldii</i> | cow clover | grassland; moist sites | perennial |
| <i>Vicia americana</i> var. <i>americana</i> | American vetch | moist woodland; forest; grassland | perennial |
| <i>Vicia gigantea</i> | giant vetch | scrub; seeps; streamsides | perennial |
| <i>Vicia lathyroides</i> * | pea vetch | disturbed sites, trail- and roadsides | annual |
| <i>Vicia sativa</i> ssp. <i>nigra</i> * | common vetch | grassland; disturbed site | annual |
| <i>Vicia tetrasperma</i> | sparrow vetch | disturbed sites | annual |

Fagaceae (Oak, Beech Family) - 9 taxa

| | | | |
|--|------------------|------------------|-------|
| <i>Chrysolepis chrysophylla</i> var. <i>chrysophylla</i> | giant chinquapin | forest | tree |
| <i>Chrysolepis chrysophylla</i> var. <i>minor</i> | dwarf chinquapin | pygmy forest | shrub |
| <i>Notholithocarpus densiflorus</i> var. <i>densiflorus</i> ¹ | tanbark oak | forest | tree |
| <i>Quercus agrifolia</i> var. <i>agrifolia</i> | coast live oak | woodland; forest | tree |
| <i>Quercus chrysolepis</i> | canyon live oak | forest | tree |
| <i>Quercus garryana</i> var. <i>garryana</i> | Oregon white oak | woodland | tree |

| | | | |
|--|-------------------------|---------------------------------------|---------------------|
| <i>Quercus kelloggii</i> | black oak | forest | tree |
| <i>Quercus parvula</i> var. <i>shrevei</i> | Shreve oak | forest | tree |
| <i>Quercus wislizeni</i> var. <i>frutescens</i> | interior live oak | forest edge | shrub |
| Garryaceae (Silttassel Family) - 1 taxon | | | |
| <i>Garrya elliptica</i> | coast silttassel | scrub; coastal bluffs | shrub |
| Gentianaceae (Gentian Family) - 3 taxa | | | |
| <i>Cicendia quadrangularis</i> | cicendia | grassland | annual |
| <i>Gentiana affinis</i> var. <i>ovata</i> | Oregon gentian | scrub | perennial |
| <i>Zeltnera davyi</i> | centaurium | moist grassland | annual |
| Geraniaceae (Geranium Family) - 7 taxa | | | |
| <i>Erodium botrys</i> * | long-beaked filaree | disturbed sites; grassland | annual |
| <i>Erodium cicutarium</i> * | red-stemmed filaree | disturbed sites; grassland | annual |
| <i>Geranium carolinianum</i> | Carolina geranium | shady woodland | annual |
| <i>Geranium core-core</i> * | New Zealand geranium | disturbed sites, roadsides | perennial |
| <i>Geranium dissectum</i> * | cutleaf geranium | disturbed sites; grassland | annual |
| <i>Geranium molle</i> * | softleaf geranium | disturbed sites; grassland | annual; biennial |
| <i>Pelargonium grossularioides</i> * | garden geranium | disturbed sites; roadsides | perennial |
| Hydrangeaceae (Hydrangea Family) - 1 taxon | | | |
| <i>Whipplea modesta</i> | modesty; yerba de selva | shaded woodland banks | subshrub |
| Hypericaceae (St. John's Wort Family) = Clusiaceae (Best, et al.) - 2 taxa | | | |
| <i>Hypericum anagalloides</i> | Tinker's penny | wet meadow | perennial |
| <i>Hypericum perforatum</i> * | Klamath weed | disturbed sites, trail- and roadsides | perennial |
| Lamiaceae (Mint Family) - 8 taxa | | | |
| <i>Clinopodium douglasii</i> | yerba buena | moist scrub understory; woodland | perennial |
| <i>Mentha pulegium</i> * | pennyroyal | moist to wet places | perennial |
| <i>Mentha spicata</i> * | spearmint | moist to wet places | perennial |
| <i>Monardella villosa</i> ssp. <i>villosa</i> | hairy coyote mint | coastal bluffs; scrub | perennial; subshrub |
| <i>Pogogyne zizyphoroides</i> | Sacramento beardstyle | wet depressions | annual |
| <i>Prunella vulgaris</i> var. <i>lanceolata</i> | self-heal | moist places; grassland | perennial |
| <i>Stachys rigida</i> var. <i>quercetorum</i> | hedge-nettle | grassland; scrub; woodland | perennial |
| <i>Stachys chamissonis</i> | marsh hedge-nettle | wet places | perennial |
| Limnanthaceae (Meadowfoam Family) - 1 taxon | | | |
| <i>Limnanthes douglasii</i> ssp. <i>douglasii</i> | Douglas meadowfoam | wet grassland; marshes | annual |
| Linaceae (Flax Family) - 2 taxa | | | |

| | | | |
|---|-------------------------------------|---------------------------------|--|
| <i>Linum bienne</i> * | blue flax | grassland | perennial |
| <i>Linum usitatissimum</i> * | common flax | disturbed sites, roadsides | annual |
| Lythraceae (Loosestrife Family) - 1 taxon | | | |
| <i>Lythrum hyssopifolia</i> * | hyssop-leaved loosestrife | moist, disturbed sites | annual; biennial |
| Malvaceae (Mallow Family) - 2 taxa | | | |
| <i>Sidalcea malviflora</i> ssp. <i>malviflora</i> | mallow-flowered (duh!) checkerbloom | grassland | perennial |
| 1B.2 <i>Sidalcea malviflora</i> ssp. <i>purpurea</i> | coastal purple checkerbloom | grassland (Gerstle Cove area) | perennial |
| Montiaceae (Montia Family) - 4 taxa | | | |
| <i>Calandrinia menziesii</i> | red maids | grassland; sandy areas | annual |
| <i>Claytonia parviflora</i> | claytonia | gravelly, sandy slopes in scrub | annual |
| <i>Claytonia perfoliata</i> | miner's lettuce | grassland; scrub; woodland | annual |
| <i>Claytonia sibirica</i> | candy flower | moist scrub | annual/perennial |
| Myricaceae (Wax Myrtle Family) - 1 taxon | | | |
| <i>Morella californica</i> | wax myrtle | moist canyons, slopes | shrub |
| Myrsinaceae - 3 taxa | | | |
| <i>Lysimachia arvensis</i> * | scarlet pimpernel | disturbed sites; grassland | annual |
| <i>Lysimachia latifolia</i> | starflower | woodland, forest | perennial |
| <i>Lysimachia minima</i> | chaffweed | wet places in grassland | annual |
| Myrtaceae (Myrtle Family) - 1 taxon | | | |
| <i>Eucalyptus globulus</i> * | Tasmanian bluegum | disturbed sites | tree |
| Onagraceae (Evening Primrose Family) - 6 taxa | | | |
| <i>Camissoniopsis cheiranthifolia</i> | beach evening-primrose | coastal strand; dunes | perennial |
| <i>Chamerion angustifolium</i> ssp. <i>circumvagum</i> | fireweed | moist, open places | perennial |
| <i>Clarkia davyi</i> | Davy clarkia | grassland | annual |
| <i>Epilobium ciliatum</i> ssp. | fringed willow-herb | wet places | perennial |
| <i>Fuchsia boliviiana</i> * | fuchsia | gardens, old homesteads | shrub |
| <i>Taraxia ovata</i> | sun cup | moist grassland | perennial |
| Orobanchaceae (Broom-rape Family) - 9 taxa | | | |
| <i>Castilleja ambigua</i> ssp. <i>ambigua</i> | johnny-nip | saline, marshy areas | annual |
| <i>Castilleja attenuata</i> | valley tassels | grassland | annual |
| <i>Castilleja latifolia</i> | Monterey Coast paintbrush | coastal bluffs | perennial, subshrub |
| <i>Castilleja subinclusa</i> ssp. <i>franciscana</i> | Franciscan paintbrush | dry slopes (Miller Cr. Canyon) | perennial |
| <i>Castilleja wightii</i> | Wight's paintbrush | scrub; coastal bluffs | perennial, subshrub |
| <i>Pedicularis densiflora</i> | ruby spires | maritime chaparral, woodland | perennial (with Arbutus or Arctostaphylos) |

| | | | |
|---|--------------------------------|------------------------------------|------------------------------|
| <i>Triphysaria eriantha</i> ssp. <i>rosea</i> | johhny-tuck | grassland | annual |
| <i>Triphysaria pusilla</i> | dwarf owl's-clover | disturbed sites; grassland | annual |
| <i>Triphysaria versicolor</i> ssp. <i>versicolor</i> | yellow-beak false owl's-clover | grassland | annual |
| Oxalidaceae (Oxalis Family) - 4 taxa | | | |
| <i>Oxalis corniculata</i> * | creepy oxalis | disturbed sites | perennial |
| <i>Oxalis oregana</i> | redwood sorrel | redwood forest | perennial |
| <i>Oxalis pes-caprae</i> * | Bermuda buttercup | disturbed sites; grassland | perennial |
| <i>Oxalis pilosa</i> | hairy wood sorrel | grassland | perennial |
| Papaveraceae (Poppy Family) - 3 taxa | | | |
| <i>Dicentra formosa</i> | western bleeding heart | moist redwood forest | perennial |
| <i>Eschscholzia californica</i> | California poppy | grassland | annual; perennial |
| <i>Platystemon californicus</i> | cream cups | grassland | annual |
| Phrymaceae (Phryma Family) - 3 taxa | | | |
| <i>Diplacus aurantiacus</i> | sticky monkeyflower | scrub; woodland | perennial; subshrub to shrub |
| <i>Erythranthe guttata</i> | seep monkeyflower | wet places | annual, perennial |
| <i>Erythranthe moschata</i> | musk monkeyflower | marshes | perennial |
| Plantaginaceae (Plantain Family) - 10 taxa | | | |
| <i>Callitricha heterophylla</i> | water-starwort | pond edges, marshes, wet places | annual |
| <i>Digitalis purpurea</i> * | foxglove | moist forest, other sites | perennial |
| <i>Gratiola ebracteata</i> | bractless hedge-hyssop | marshy, muddy sites | annual |
| <i>Kickxia spuria</i> * | fluellin | disturbed sites | annual |
| <i>Plantago coronopus</i> * | cutleaf plantain | disturbed sites; grassland; bluffs | annual; biennial |
| <i>Plantago erecta</i> | Californiaplantain | grassland | annual |
| <i>Plantago lanceolata</i> * | English plantain | disturbed sites; grassland | perennial |
| <i>Plantago major</i> * | common plantain | disturbed moist sites | perennial |
| <i>Plantago maritima</i> | seaside plantain | coastal bluffs | perennial |
| <i>Plantago subnuda</i> | almost-naked plantain | marshes; moist bluffs, banks | perennial |
| <i>Synthyris reniformis</i> | snow queen | moist, shady sites; woodland | perennial |
| Plumbaginaceae (Leadwort Family) - 2 taxa | | | |
| <i>Armeria maritima</i> ssp. <i>californica</i> | sea thrift; sea-pink | coastal bluffs; coastal grassland | perennial |
| <i>Limonium californicum</i> | sea-lavender; marsh-rosemary | salt marsh; moist bluffs | perennial |
| Polemoniaceae (Phlox Family) - 5 taxa | | | |
| <i>Collomia heterophylla</i> ¹ | vari-leaf collomia | moist scrub; woodland | annual |
| 1B.1 <i>Gilia capitata</i> ssp. <i>tomentosa</i> | tomentose globe gilia | Miller Canyon | annual |

| | | | |
|--|-------------------------|----------------------------------|------------------|
| <i>Leptosiphon bicolor</i> | baby stars | grassland | annual |
| <i>Navarretia mellita</i> | honey-scented skunkweed | sandy slopes (Stump Beach) | annual |
| <i>Navarretia squarrosa</i> | skunkweed | grassland; disturbed sites | annual |
| Polygalaceae (Milkwort Family) - 1 taxon | | | |
| <i>Polygala californica</i> | California milkwort | dry woodland | perennial |
| Polygonaceae (Knotweed Family) - 8 taxa | | | |
| <i>Chorizanthe cuspidata</i> var. <i>villosa</i> | spineflower | coastal, sandy soils | annual |
| <i>Eriogonum latifolium</i> | coast buckwheat | grassland, coastal bluffs | perennial |
| <i>Polygonum aviculare</i> ssp. <i>depressum</i> * | dooryard knotweed | disturbed sites | annual |
| <i>Polygonum paronychia</i> | dune knotweed | coastal strand; grassland | subshrub |
| <i>Rumex acetosella</i> * | sheep sorrel | disturbed sites, grassland | perennial |
| <i>Rumex conglomeratus</i> * | s | low-lying, moist areas | perennial |
| <i>Rumex crispus</i> * | curly dock | disturbed moist sites | perennial |
| <i>Rumex crassus</i> | willow dock | sandy coastal sites; dunes | perennial |
| <i>Rumex obtusifolius</i> * | bitter dock | shaded, moist openings in forest | perennial |
| Ranunculaceae (Buttercup Family) - 4 taxa | | | |
| <i>Actaea rubra</i> | baneberry | moist forest slopes | perennial |
| <i>Aquilegia formosa</i> | western columbine | moist sites | perennial |
| <i>Ranunculus californicus</i> | California buttercup | moist grassland | perennial |
| <i>Ranunculus muricatus</i> * | stickseed buttercup | moist, disturbed sites | annual; biennial |
| Rhamnaceae (Buckthorn Family) - 8 taxa | | | |
| <i>Ceanothus foliosus</i> var. <i>foliosus</i> | wavyleaf ceanothus | dry woodland; scrub | shrub |
| 4.3 <i>Ceanothus gloriosus</i> var. <i>exaltatus</i> | glory brush | upper praire | |
| 4.3 <i>Ceanothus gloriosus</i> var. <i>gloriosus</i> | Pt. Reyes ceanothus | coastal bluffs, slopes | shrub |
| <i>Ceanothus thyrsiflorus</i> var. <i>griseus</i> | Carmel ceanothus | coastal scrub, slopes | shrub |
| <i>Ceanothus thyrsiflorus</i> var. <i>thyrsiflorus</i> | blue blossom | woodland; scrub | shrub |
| <i>Ceanothus velutinus</i> | tobacco brush | woodland slopes | shrub |
| <i>Frangula californica</i> | coffeberry | woodland; scrub | shrub |
| <i>Frangula purshiana</i> | cascara sagrada | moist woodland | shrub |
| Rosaceae (Rose Family) - 22 taxa | | | |
| <i>Acaena novae-zelandiae</i> * | biddy-biddy | grassland | perennial |
| <i>Acaena pinnatifida</i> | acaena | grassland | perennial |
| <i>Amelanchier alnifolia</i> | service-berry | woodland; open slopes; scrub | shrub |
| <i>Aphanes occidentalis</i> | lady's mantle | disturbed sites; grassland | annual |

| | | | |
|--|------------------------|--------------------------------|----------------------|
| <i>Cotoneaster lacteus</i> * | Parney cotoneaster | disturbed sites, grassland | shrub |
| <i>Cotoneaster pannosus</i> * | silverleaf cotoneaster | scrub; disturbed sites | shrub |
| <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> | sticky cinquefoil | grassland; woodland | perennial |
| <i>Fragaria chiloensis</i> | beach strawberry | beach; dunes; grassland; scrub | perennial (trailing) |
| <i>Fragaria vesca</i> | wood strawberry | scrub; grassland; woodland | perennial (trailing) |
| <i>Heteromeles arbutifolia</i> | toyon | scrub; woodland | shrub |
| <i>Horkelia californica</i> | California horkelia | scrub; grassland | perennial |
| 1B.2 <i>Horkelia tenuiloba</i> | thin-lobed horkelia | grassland | perennial |
| <i>Potentilla anserina</i> ssp. <i>pacifica</i> | silverweed | wet coastal places | perennial |
| <i>Pyracantha angustifolia</i> * | firethorn | disturbed sites | shrub |
| <i>Rosa californica</i> | California wild rose | scrub; moist places | shrub |
| <i>Rosa rubiginosa</i> * | sweet-brier | scrub; disturbed places | shrub |
| <i>Rosa gymnocarpa</i> | wood rose | woodland; forest | perennial; shrub |
| <i>Rosa nutkana</i> var. <i>nutkana</i> | Nootka rosa | moist flat areas | shrub |
| <i>Rubus armeniacus</i> * | Armenian blackberry | disturbed sites | shrub |
| <i>Rubus leucodermis</i> | blackcap raspberry | slopes; canyons | shrub |
| <i>Rubus parviflorus</i> | thimbleberry | moist places; scrub | shrub |
| <i>Rubus spectabilis</i> | salmonberry | moist places; streamsides | shrub |
| <i>Rubus ursinus</i> | California blackberry | grassland; scrub | shrub; trailing |

Rubiaceae (Madder Family) - 6 taxa

| | | | |
|--|---------------------------------|--|----------------------|
| <i>Galium aparine</i> | goose-grass; bedstraw; cleavers | grassland; woodland | annual (trailing) |
| <i>Galium californicum</i> ssp. <i>maritimum</i> | bedstraw | woodland; forest | perennial (trailing) |
| <i>Galium muricatum</i> | Oregon bedstraw | forest (Praire Trail nr. Plantation Rd.) | perennia |
| <i>Galium porrigens</i> var. <i>porrigens</i> | climbing bedstraw | woodland; scrub; forest | perennial (climbing) |
| <i>Galium trifidum</i> var. <i>pacificum</i> | bedstraw | wet places | perennial |
| <i>Galium triflorum</i> | sweet-scented bedstraw | woodland | perennial |
| <i>Sherardia arvensis</i> * | field madder | grassland | annual |

Salicaceae (Willow Family) - 5 taxa

| | | | |
|--|------------------|-------------------|-------------|
| <i>Salix hookeriana</i> | coastal willow | stream mouths | shrub |
| <i>Salix laevigata</i> | red willow | streamsides | shrub; tree |
| <i>Salix lasiandra</i> ssp. <i>lasiandra</i> | shining willow | wet places; seeps | shrub; tree |
| <i>Salix scouleriana</i> | Scouler's willow | woodland | tree |
| <i>Salix sitchensis</i> | Sitka willow | streamsides | shrub; tree |

Sapindaceae (Soapberry Family)) -- 1 taxon

| | | | |
|--|---------------------------|--|--------------------------------|
| <i>Aesculus californica</i> | California buckeye | woodland; slopes | tree |
| Saxifragaceae (Saxifrage Family) - 5 taxa | | | |
| <i>Boykinia occidentalis</i> | western boykinia | creek banks | perennial |
| <i>Heuchera micrantha</i> | alumroot | damp banks, rocks, cliffs | perennial |
| <i>Heuchera pilosissima</i> | coast alumroot | coastal bluffs, slopes | perennial |
| <i>Tiarella trifoliata</i> var. <i>unifoliata</i> | sugar-scoop | moist, shaded slopes | perennial |
| <i>Tolmiea diplomenziesii</i> | pig-a-back plant | damp, cool places | perennial |
| Scrophulariaceae (Figwort Family) - 1 taxon | | | |
| <i>Scrophularia californica</i> ssp. <i>californica</i> | bee plant | moist to wet places; scrub | perennial |
| Solanaceae (Nightshade Family) - 2 taxa | | | |
| <i>Solanum americanum</i> | small-flowered nightshade | moist places | annual, perennial, subshrub |
| <i>Solanum furcatum</i> * | forked nightshade | disturbed sites, bluffs at Stump Beach | perennial |
| Urticaceae (Nettle Family) - 1 taxon | | | |
| <i>Urtica dioica</i> ssp. <i>gracilis</i> | American stinging nettle | wet thickets; streamsides | perennial |
| Violaceae (Violet Family) - 3 taxa | | | |
| <i>Viola adunca</i> | western dog violet | grassland | perennial |
| <i>Viola glabella</i> | stream violet | along streams | perennial |
| <i>Viola sempervirens</i> | redwood violet | woodland, forest | perennial |
| Viscaceae (Mistletoe Family) - 1 taxon | | | |
| <i>Arceuthobium campylopodum</i> | western dwarf mistletoe | on <i>Abies</i> , <i>Pinus</i> | perennial (hemi-parasite) |
| MONOCOTS | | | |
| Agavaceae (Yucca Family) - 3 taxa | | | |
| <i>Camassia quamash</i> ssp. <i>breviflora</i> | common camas | moist grassland; marsh | perennial |
| <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i> | soap root | woodland | perennial |
| <i>Cordyline australis</i> * | cordyline | forest; riparian | perennial (shrub/tree) |
| Alliaceae (Onion Family) - 2 taxa | | | |
| <i>Allium dichlamydeum</i> | coast onion | grassland; rocky bluffs | perennial (bulb) |
| <i>Allium unifolium</i> | single-leaf onion | grassland; open sites | perennial (bulb) |
| Amaryllidaceae (Amaryllis Family) - 1 taxon | | | |
| <i>Narcissus pseudonarcissus</i> * | daffodil | disturbed sites (garden escape) | perennial (bulb) |
| Araceae (Arum Family) - 2 taxa | | | |
| <i>Lemna minor</i> | duckweed | surface of pools, ponds, slow-moving water | perennial |
| <i>Zantedeschia aethiopica</i> * | calla lily | seeps; streamsides; springs; moist grassland | perennial |

| | | | | |
|--|---|----------------------|--|---------------------|
| Asphodelaceae (Asphodel Family) - 1 taxon | | | | |
| | <i>Kniphofia uvaria</i> * | red-hot poker | disturbed sites | perennial (rhizome) |
| Cyperaceae (Sedge Family) - 16 taxa | | | | |
| | <i>Carex bolanderi</i> | Bolander sedge | ditches, marshes | perennial |
| 2.3 | <i>Carex californica</i> | California sedge | woodland, trails | perennial |
| | <i>Carex globosa</i> | round-fruited sedge | woodland | perennial |
| | <i>Carex gynodynama</i> | wonder woman sedge | moist grassland, trails | perennial |
| | <i>Carex harfordii</i> | Harford sedge | moist grassland, forest openings | perennial |
| | <i>Carex hendersonii</i> | Henderson sedge | shaded, moist woodland, forest edge | perennial |
| | <i>Carex leptopoda</i> | slender-footed sedge | redwood forest | perennial |
| | <i>Carex obnupta</i> | slough sedge | marshes; wet grassland, scrub; seeps | perennial |
| | <i>Carex tumulicola</i> | foothill sedge | open, seasonally moist grassland, woodland | perennial |
| | <i>Carex vesicaria</i> | inflated sedge | marshes; wet grassland | perennial |
| | <i>Cyperus eragrostis</i> var. <i>eragrostis</i> | love-grass nut-sedge | moist grassland, woodland, disturbed sites | perennial |
| | <i>Isolepis carinata</i> | bulrush | wet ground; seeps | perennial |
| | <i>Isolepis cernua</i> | low lateral bulrush | wet ground; marshes; seeps | perennial |
| | <i>Schoenoplectus americanus</i> | American bulrush | marshes; streamsides | perennial |
| | <i>Schoenoplectus pungens</i> var. <i>longispicatus</i> | threesquare | marshes; wet, alkaline soils | perennial |
| | <i>Scirpus microcarpus</i> | panicked bulrush | seeps; streamsides; marshes | perennial |
| Iridaceae (Iris Family) - 7 taxa | | | | |
| | <i>Crocosmia X crocosmiiflora</i> * | montbretia | disturbed sites, roadsides | perennial (corm) |
| | <i>Iris douglasiana</i> | Douglas iris | grassland; open woodland | perennial (rhizome) |
| 4.2 | <i>Iris longipetala</i> | coast iris | moist, open sites | perennial (rhizome) |
| | <i>Romulea rosea</i> ssp. <i>australis</i> * | rosy sand crocus | disturbed roadside gravel, grassland | perennial (corm) |
| | <i>Sisyrinchium bellum</i> | blue-eyed grass | grassland; open woodland | perennial (rhizome) |
| | <i>Sisyrinchium californicum</i> | golden-eyed grass | moist to wet, open sites | perennial (rhizome) |
| | <i>Watsonia meriana</i> * | watsonia | disturbed sites (garden escape) | perennial (corm) |
| Juncaceae (Rush Family) - 12 taxa | | | | |
| | <i>Juncus bolanderi</i> | Bolander's rush | wet ground; marshes; seeps | perennial |
| | <i>Juncus bufonius</i> var. <i>bufonius</i> | toad rush | seasonally moist ground | annual |
| | <i>Juncus capitatus</i> * | rush | vernally wet grassland, slopes | annual |
| | <i>Juncus effusus</i> ssp. <i>pacificus</i> | Pacific rush | marshes; wet grassland, seeps, slopes | perennial |
| | <i>Juncus ensifolius</i> | rush | marshes; wet grassland | perennial |
| | <i>Juncus hesperius</i> | coast rush | ditches; marshes | perennial |

| | | | |
|---|--------------------|---------------------------------------|-----------|
| <i>Juncus lescurii</i> | San Francisco rush | sandy flats, slopes, grassland; dunes | perennial |
| <i>Juncus occidentalis</i> | western rush | seasonally moist ground | perennial |
| <i>Juncus patens</i> | spreading rush | damp grassland, slopes; streamsides | perennial |
| <i>Juncus phaeocephalus</i> var. <i>phaeocephalus</i> | brown-headed rush | marshes; wet ground | perennial |
| <i>Juncus xiphoides</i> | iris-leaved rush | marshes; springs | perennial |
| <i>Luzula comosa</i> | wood rush | moist grassland, woodland, slopes | perennial |

Liliaceae (Lily Family) - 9 taxa; see also Agavaceae, Alliaceae, Amaryllidaceae, Asphodelaceae, Melanthiaceae, Ruscaceae, Themidaceae)

| | | | | |
|-------------|---|----------------------|--|---------------------|
| | <i>Calochortus tolmiei</i> | pussy ears | grassland; woodland | perennial (bulb) |
| 4.2 | <i>Calochortus uniflorus</i> | pink star-tulip | grassland; woodland | perennial (bulb) |
| | <i>Clintonia andrewsiana</i> | blue bead lily | moist, shaded woodland | perennial (rhizome) |
| | <i>Lilium columbianum</i> | Oregon lily | dry scrub, grassland, forest openings | perennial (rhizome) |
| 1B.1 | <i>Lilium maritimum</i> | coast lily | moist, sandy sites in woodland, forest | perennial (rhizome) |
| | <i>Lilium pardalinum</i> ssp. <i>pardalinum</i> | leopard lily | springs, seeps, streamsides | perennial (rhizome) |
| | <i>Prosartes hookeri</i> | Hooker fairy bells | shaded sites; woodland; forest | perennial (rhizome) |
| | <i>Prosartes smithii</i> | Smith fairy bells | coastal shaded, moist sites (Miller Cr.) | perennial (rhizome) |
| | <i>Scoliopus bigelovii</i> | fetid adder's tongue | moist, shaded woodland, forest | perennial (rhizome) |

Melanthiaceae (False-hellebore Family) 5 taxa

| | | | | |
|------------|---|-------------------------|---|---------------------|
| | <i>Toxicoscordion fontanum</i> | star lily | moist sites; flats | perennial (rhizome) |
| | <i>Toxicoscordion fremontii</i> | Fremont death-camas | drier sites, slopes; scrub; woodland; grassland | perennial (rhizome) |
| | <i>Trillium ovatum</i> ssp. <i>ovatum</i> | wake robin | moist woods, scrub | perennial (rhizome) |
| 4.3 | <i>Veratrum fimbriatum</i> | fringed false hellebore | wet sites; streamsides; meadows | perennial (rhizome) |
| | <i>Xerophyllum tenax</i> | bear-grass | dry, open sites | perennial (rhizome) |

Orchidaceae (Orchid Family) - 7 taxa

| | | | | |
|--|----------------------------------|----------------------|--|--------------------------------|
| | <i>Calypso bulbosa</i> | calypso orchid | shaded forest | perennial |
| | <i>Corallorrhiza maculata</i> | spotted coralroot | shaded forest | perennial |
| | <i>Corallorrhiza mertensiana</i> | western coralroot | shaded forest, in humus | perennial (non-green, rhizome) |
| | <i>Corallorrhiza striata</i> | striped coralroot | dense, shaded woodland, forest | perennial (non-green, rhizome) |
| | <i>Epipactis gigantea</i> | stream orchid | streamsides | perennial (rhizome) |
| | <i>Epipactis helleborine</i> * | broad-leaved orchid | forest (Praire Trail nr. Plantation Rd.) | perennial (rhizome) |
| | <i>Goodyera oblongifolia</i> | rattlesnake-plantain | shaded forest | perennial |
| | <i>Spiranthes romanzoffiana</i> | ladies' tresses | moist grassland, seeps | perennial (rhizome) |

Poaceae (Grass Family) aka Graminae - 61 taxa

| | | | | |
|--|-------------------------|-----------------|---------------------|----------------------|
| | <i>Agrostis exarata</i> | spikebentgrass | woodland | perennial |
| | <i>Agrostis pallens</i> | thin bent grass | grassland; woodland | perennial (rhizomes) |

| | | | | |
|-----|--|---------------------------|---|-----------------------|
| | <i>Agrostis stolonifera</i> * | redtop bentgrass | moist areas | perennial |
| | <i>Aira caryophyllea</i> * | silver European hairgrass | grassland; disturbed sites | annual |
| | <i>Aira praecox</i> * | hairgrass | bluffs; grassland | annual |
| | <i>Ammophila arenaria</i> * | European beachgrass | dunes; beaches | perennial (rhizome) |
| | <i>Anthoxanthum occidentale</i> | vanilla grass | forest; scrub | perennial (rhizomes) |
| | <i>Anthoxanthum odoratum</i> * | sweet vernal grass | moist grassland | perennial (cespitate) |
| | <i>Arundo donax</i> * | giant reed | seeps; wetlands (Stump Beach) | perennial (rhizomes) |
| | <i>Avena sativa</i> * | cultivated oat | disturbed sites | annual |
| | <i>Briza maxima</i> * | rattlesnake grass | grassland; disturbed sites; roadcuts | annual |
| | <i>Briza minor</i> * | little quaking grass | grassland; disturbed sites | annual |
| | <i>Bromus carinatus</i> var. <i>carinatus</i> | California brome | grassland; woodland | perennial (cespitate) |
| | <i>Bromus carinatus</i> var. <i>maritimus</i> | California brome | dunes; coastal bluffs, grassland | perennial (cespitate) |
| | <i>Bromus diandrus</i> * | ripgut brome | sandy soils; grassland; disturbed sites | annual |
| | <i>Bromus hordeaceus</i> * | soft chess | grassland; woodland; disturbed sites | annual |
| | <i>Bromus laevipes</i> | woodland brome | woodlands, forest openings | perennial |
| | <i>Bromus madritensis</i> ssp. <i>rubens</i> * | red brome | grassland; disturbed sites | annual |
| 4.2 | <i>Calamagrostis bolanderi</i> | Bolander's reed grass | forest edges, woodland | perennial (rhizome) |
| | <i>Calamagrostis nutkaensis</i> | Pacific reed grass | moist coastal grassland | perennial (cespitate) |
| | <i>Cortaderia jubata</i> * | jubatagrass | disturbed sites; bluffs; roadcuts | perennial (cespitate) |
| | <i>Cynodon dactylon</i> * | Bermuda grass | disturbed sites | perennial (rhizomes) |
| | <i>Cynosurus echinatus</i> * | hedgehog dogtail | grassland; openings in woodlands | annual |
| | <i>Dactylis glomerata</i> * | orchard grass | grassland; disturbed sites | perennial (rhizomes) |
| | <i>Danthonia californica</i> | California oatgrass | moist grassland | perennial (cespitate) |
| | <i>Deschampsia cespitosa</i> ssp. <i>holciformis</i> | tufted hairgrass | grassland | perennial (cespitate) |
| | <i>Deschampsia elongata</i> | slender hairgrass | moist grassland, openings in woodlands | perennial (cespitate) |
| | <i>Digitaria ischaemum</i> * | crabgrass | disturbed sites | annual |
| | <i>Distichlis spicata</i> | saltgrass | salt marshes, saline seeps | perennial (rhizomes) |
| | <i>Elymus glaucus</i> | blue wildrye | grassland; woodland | perennial (rhizomes) |
| | <i>Elymus pacificus</i> | Pacific wildrye | coastal bluffs, beach | perennial (rhizomes) |
| | <i>Elymus trachycaulus</i> ssp. <i>trachycaulis</i> | bearded wheatgrass | moist grassland, woodland | perennial (rhizomes) |
| | <i>Festuca arundinacea</i> * | tall fescue | moist grassland; disturbed sites | perennial (cespitate) |
| | <i>Festuca bromoides</i> * | brome fescue | moist areas; marshy ground | annual |
| | <i>Festuca californica</i> | California fescue | scrub; grassland; woodland | perennial (cespitate) |
| | <i>Festuca idahoensis</i> | Idaho fescue | scrub; grassland; woodland | perennial (cespitate) |

| | | | |
|--|----------------------------|---|-----------------------|
| <i>Festuca myuros</i> * | 6-weeks fescue | many habitats | annual |
| <i>Festuca occidentalis</i> | western fescue | forest openings | perennial (cespítose) |
| <i>Festuca perennis</i> * | Italian/perennial ryegrass | grassland, disturbed sites | annual/perennial |
| <i>Festuca pratensis</i> * | meadow fescue | disturbed sites; roadsides | perennial (cespítose) |
| <i>Festuca rubra</i> | red fescue | grassland | perennial (cespítose) |
| <i>Holcus lanatus</i> * | purple velvet grass | moist grassland; disturbed sites | perennial (clumping) |
| <i>Hordeum brachyantherum</i> ssp. <i>brachyantherum</i> | meadow barley | moist grassland | perennial (rhizomes) |
| <i>Hordeum brachyantherum</i> ssp. <i>californicum</i> | California barley | scrub; grassland | perennial (rhizomes) |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> | Mediterranean barley | grassland | annual |
| <i>Hordeum murinum</i> * ssp. <i>leporinum</i> | hare barley | grassland; disturbed sites | annual |
| <i>Lagurus ovatus</i> * | hare's tail | grassland; disturbed sites | annual |
| <i>Melica torreyana</i> | Torrey melic | woodland; scrub; forest | perennial (rhizomes) |
| <i>Panicum acuminatum</i> var. <i>fasciculatum</i> | Pacific panic grass | moist grassland (upper prairie), roadside ditches | perennial |
| <i>Phalaris aquatica</i> * | Harding grass | moist, disturbed sites | perennial (clumped) |
| <i>Phalaris californica</i> | California canary grass | marshes, roadside ditches | perennial |
| <i>Phleum pratense</i> * | cultivated timothy | disturbed sites | perennial (clumped) |
| <i>Poa annua</i> * | annual bluegrass | moist, disturbed sites | annual |
| <i>Poa douglasii</i> | sand dune bluegrass | dunes; beaches | perennial (rhizomes) |
| <i>Poa kelloggii</i> | Kellogg bluegrass | partly shaded redwood, bishop pine forest | perennial (rhizome) |
| <i>Poa unilateralis</i> | ocean bluff bluegrass | grassland; coastal bluffs, rock outcrops | perennial (cespítose) |
| <i>Polypogon interruptus</i> * | ditch beard grass | low-lying damp places | perennial (clumped) |
| <i>Polypogon monspeliensis</i> * | rabbitfoot grass | low-lying, vernally wet places | annual |
| <i>Rytidosperma penicillatum</i> * | hairy oatgrass | disturbed sites | perennial (cespítose) |
| <i>Stipa lepida</i> | foothill needlegrass | scrub; grassland | perennial (cespítose) |
| <i>Stipa pulchra</i> | purple needlegrass | grassland | perennial (cespítose) |
| <i>Trisetum canescens</i> | tall false oat | moist areas | perennial (cespítose) |
| Ruscaceae (Butcher's-groom Family) 2 taxa | | | |
| <i>Maianthemum racemosum</i> | false Solomon's-seal | woodland, forest | perennial (rhizome) |
| <i>Maianthemum stellatum</i> | false Solomon's-seal | moist sites; scrub; woodland; forest | perennial (rhizome) |
| Themidaceae (Brodiaea Family) 5 taxa | | | |
| <i>Brodiaea terrestris</i> ssp. <i>terrestris</i> | ground brodiaea | grassland; open woodland | perennial (corm) |
| <i>Dichelostemma capitatum</i> ssp. <i>capitatum</i> | blue dicks | grassland; open sites, banks | perennial (corm) |
| <i>Dichelostemma congesta</i> | ookow | woodland slopes | perennial (corm) |
| <i>Triteleia laxa</i> | Ithuriel's spear | dry flats, slopes in scrub, woodland | perennial (corm) |

| | | | |
|--|---------------------------------|-------------------------|---------------------|
| <i>Triteleia peduncularis</i> | long-rayed brodiaea | wet grassland; marshes | perennial (corm) |
| Typhaceae (Cattail Family) including Sparganiaceae (Best, et al.) - 2 taxa | | | |
| <i>Sparganium erectum</i> ssp. <i>stoloniferum</i> | bur-reed | ponds, marshes, streams | perennial (rhizome) |
| <i>Typha latifolia</i> | soft-flag; broad-leaved cattail | pond edges, marshes | perennial (rhizome) |

Notes:

1. Taxon not documented as occurring specifically at Salt Point, Horseshoe Cove, Gerstle Cove, Stump Beach in Best, et al., A Flora of Sonoma County (1996).
 2. Nomenclature or spelling according to Best, et al. (1996)
 3. Nomenclature according to Patterson, et al. (Madroño 2002)
- α left column: section/group of vascular plants (as in Jepson Manual, 2nd ed.), plant family, and where applicable, California Rare Plant Rank

Total: ~ 536 taxa

Documentation of taxa according to Best, et al. (Flora of Sonoma County, 1996), and numerous personal observations by many contributors

Taxonomic nomenclature according to the Jepson Herbarium: Jepson Flora Project: Jepson eFlora, 06/01/2015

Numbers in left hand column indicate State Rare Rank

List edited 08/02/2019 by P. Warner